

Toxic Drug Crisis Events and Deaths and FNHA's Response COMMUNITY SITUATION REPORT: MAY 2022

FNHA Public Health Response Last updated: October 6, 2022

Introduction

Each month, the First Nations Health Authority (FNHA) reports on the number of toxic drug poisoning events and deaths that have taken place among First Nations populations in BC. In the report, the FNHA also summarizes the actions that the FNHA is taking in response to the toxic drug emergency. This report covers the period January 1, 2021 to May 31, 2022. For previous reports, see FNHA's harm reduction webpage.

Summary Update (May 2022)

First Nations Toxic Drug Poisoning Events and Deaths

In May 2022, there were a total of 252 paramedic-attended drug poisoning events reported among First Nations people. This represents a 2% increase from the previous month and a 16.8% decrease from May of last year.

First Nations people represented 17% of all toxic drug poisoning events this month.

Women represented 35.8% of all First Nations toxic drug poisoning events; among other residents, 23.3% of all drug poisoning events were women.

In May, we lost 27 First Nations people due to toxic drug poisoning. First Nations people represented 14% of all deaths this month. Since 2016, the year in which a public health emergency was declared, we have lost 1,353 First Nations people to toxic drug poisoning.

FNHA's Response to the Toxic Drug Emergency

As described in the FNHA Programs and Outcomes section of this report, the FNHA has developed an expanding range of programs and initiatives to combat the toxic drug crisis. These are designed in culturally safe ways that confront the anti-Indigenous racism and systemic inequity built into Canada's health system.

Key programs include First Nations Treatment and Healing Centres, Intensive Case Management (ICM) Teams, Indigenous land-based healing services, Not Just Naloxone training, the development of a network of peer coordinators, hiring of community-facing harm reduction educators, dispensing opioid agonist therapy (OAT), and distributing naloxone.

Provision of OAT

Based on prescription drug claim data of FNHA clients, 2,557 First Nations people were dispensed OAT in May 2022. Of these:



- 60.4% were dispensed methadone, 22.1% were dispensed buprenorphine/naloxone (Suboxone), 16.9% were dispensed slow-release oral morphine (Kadian) and a small percent were dispensed buprenorphine extended-release (Sublocade)
- 2.5% were dispensed OAT through FNHA Health Benefits for the first time


Naloxone Distribution

- Through FNHA's bulk ordering program, 1060 nasal naloxone kits were distributed to First
 Nations and community organizations in May 2022 (each kit contains two doses), and another
 193 doses of nasal naloxone were distributed to FNHA clients through community pharmacies¹
- 720 injectable naloxone kits were ordered for First Nations sites or Friendship Centres (these kits contain three doses)

Toxic Drug Poisoning Events and Deaths Data (January 1, 2021 to May 31, 2022)

Paramedic-Attended Events by Month


Since COVID-19 was declared a pandemic, there have been increases in the number of both toxic drug events and deaths among First Nations people, although there has been a decline in drug poisoning events since the peak in July 2021.



¹ Effective November 26, 2021, First Nations individuals in BC may obtain 2 kits (4 cartridges) of nasal naloxone every 30 days from pharmacies. First Nations and community organizations that provide harm reduction support primarily to First Nations people may request bulk supply of nasal naloxone; this may be ordered along with other culturally safe harm reduction supplies through FNHA's harm reduction hub: HarmReduction@fnha.ca.



Deaths by Month



Note: Suppressed when the number of deaths is less than 10 or to avoid back-calculation of another number that is less than 10

Events and Deaths by Region (January 1, 2021 – May 31, 2022)

	Fraser Salish	Interior	Northern	Vancouver Coastal	Vancouver Island	ВС
Total Paramedic-Attended Drug Poisoning Events	919	686	903	1511	825	4844
Total Number of Deaths	78	85	86	155	82	486
Percentage of the Population that is First Nations ²	1.4%	4.1%	14.8%	2.1%	4.2%	3.3%
Percentage of all Events that were First Nations ³	9.9%	16%	51.2%	19.7%	20.7%	18.0%
Percentage of all Deaths that were First Nations*	7.2%	16.1%	38.4%	17.4%	16.8%	15.1%
Crude Drug Poisoning Event Rate (per 1,000) ⁴	84.1	49.2	52.2	149.9	55.4	71.6
OAT Claimants (in May 2022) ⁵	705	348	347	736	520	2557

^{*} The number of deaths by region and the proportion of all deaths that were First Nations are updated quarterly in order to protect privacy.

² Based on 2018 estimates from First Nations Client File (FNCF) 2018 and BC Stats Population Estimates.


³ Based on records with a complete Personal Health Number (PHN) only.

⁴ Estimated rate for 2021-2022 based on 17 months of data; 2019 population estimates via 2018 FNCF.

⁵ If a person was a claimant in two or more different regions in any given month they will count as a claimant for each region; hence, the sum of the regions is greater than the BC number presented in the table.




Paramedic-Attended Events by Age Group



Approximately 60% of all First Nations persons who had a paramedic attended drug-poisoning event in the first 5 months of 2022 were younger than 40 years of age.

Paramedic Attended Events by Sex



Note: Data on toxic drug deaths by sex is updated quarterly in order to protect privacy.

Women continue to represent higher proportions of First Nations toxic drug poisoning events and deaths compared to Other Residents.

- 35.8%% of toxic drug poisoning events among First Nations involved women, this compares to 23.3% among other residents of BC.
- 37.8% of toxic drug poisoning deaths among First Nations involved women, this compares to 21.0% among other residents of BC.



For provincial-level data, please see:


- Illicit Drug Toxicity Deaths in BC (BC Coroners Service)
- <u>Unregulated Drug Poisoning Emergency Dashboard</u> (BCCDC)

Paramedic-Attended Events by Local Health Area

The local health areas with the highest drug poisoning event rates (indicated on the map below) in the most recent 1 year (June 2021 – May 2022) were:

- Vancouver Centre North (135.3 per 1,000)
- Vancouver City Centre (106.5 per 1,000)
- Abbotsford (65.2 per 1,000)
- Vancouver Midtown (53.4 per 1,000)
- Penticton (43.3 per 1,000)
- Prince George (43 per 1,000)

The local health areas with the highest drug poisoning counts (not displayed on map) were Vancouver - Centre North, Prince George, Surrey, Kamloops, Terrace and Greater Victoria.



Note: LHAs with the highest **rates** or highest **number** of events have been labelled in the map above.



FNHA's Response to the Toxic Drug Emergency

FNHA's Toxic Drug Emergency Response Framework for Action details an iterative approach to evolving our response to the crisis based on what we hear from community members, health directors, leaders, frontline staff, peers and others throughout the process of implementation.

SYSTEM-WIDE TOXIC DRUG PUBLIC HEALTH EMERGENCY RESPONSE FOR FIRST NATIONS IN BC

4

- Prevent people who overdose from dying
- Keep people safe when using substances
- · Create an accessible range of treatment options
- Support people on their healing journeys

The full Framework is available here: <u>A Framework for Action: Responding to the Toxic Drug Crisis for</u> First Nations.

FNHA Programs and Outcomes

As the drug toxicity emergency has unfolded and worsened during the COVID-19 pandemic, the FNHA has implemented numerous ongoing and new programs and initiatives, including:

- Currently, eight First Nations treatment and healing centres operate across BC and two new facilities are being planned – one in the Vancouver Coastal region and the other in the Fraser Salish region
- Intensive Case Management (ICM) Teams provide wrap-around support for individual and family wellness and access to care in all five regions
- Indigenous land-based healing services grounded in cultural teachings are provided at 147 sites across BC
- Virtual and in-person harm reduction education through Not Just Naloxone training and community visits: from June 1, 2021 to May 31, 2022, 27 training sessions were held; 752 health care workers, youth, Elders and community champions were trained
- Broadened access to nasal spray naloxone through bulk supply ordering by First Nations communities and organizations across BC (see table below)
- Unlocking the Gates supports people who are leaving prison and are at a dramatically higher risk of overdose from toxic drugs
- Expanding the regional overdose response capacity with **new hires of harm reduction educators** and peer coordinators being deployed in urban hotspots, based on health surveillance data
- Increasing access to OAT:
 - directly through nurse prescribing: approved at five sites and in approval process at seven other sites; 35 nurses are enrolled in or have completed prescribing training
 - by supporting 30 rural and remote First Nations communities to improve access to OAT for their members




- Developed the Indigenous Harm Reduction Community Council a province-wide network of Indigenous people working on Indigenous approaches to harm reduction. The Council is coordinated by 14 members representing all five regions; a web portal for the network is under development
- Approved a <u>Harm Reduction Policy</u> with five areas for action:
 - increase access to cultural activities
 - expand access to substitution therapies (such as OAT)
 - o provide harm reduction services and promote expansion of related strategies
 - engage with people with lived and living experience in design and implementation
 - o support expansion of pharmaceutical alternatives to toxic street drugs

The FNHA also has several new and emerging initiatives:

- The FNHA is establishing three Indigenous-focused overdose prevention sites (OPS); in the
 Fraser Salish region, the FNHA partnering with Cheam First Nation and Fraser Health to
 implement a first of its kind OPS in the Cheam First Nation community
- The FNHA has coordinated with the Western Aboriginal Harm Reduction Society (WAHRS) to open an **episodic OPS** in the Downtown Eastside and is working on identifying other sites in BC for these projects, to be known as Raven's Eye Sage Sites
- The FNHA will engage with First Nations families and communities to explore the
 decriminalization of people who use substances; guided by these conversations the FNHA will
 work with system partners to ensure First Nations priorities, perspectives and experiences
 influence discussions and decisions on decriminalization in BC
- The FNHA will also engage with communities to assess the need and preferences for pharmaceutical alternative to toxic street drugs by First Nations people who are at risk of overdose

Access to OAT

Number of FNHA Clients Dispensed OAT




OAT is one of the recommended pharmacotherapy options to reduce opioid-use related harms and to support long-term recovery for persons with opioid use disorder. The medications include but are not limited to methadone, buprenorphine/naloxone (Suboxone), slow-release oral morphine (Kadian) and buprenorphine extended-release (Sublocade).




With the expansion of OAT initiatives throughout the province, the total number of FNHA clients who were dispensed any type of OAT covered by the FNHA pharmacy benefit plan has steadily increased to 2,557 persons in May 2022.

Methadone was the most commonly prescribed type of OAT among FNHA clients dispensed OAT in May 2022. 60.4% of FNHA clients dispensed any type of OAT under the FNHA health benefit plan in May 2022. were prescribed methadone, while 22.1% of were prescribed buprenorphine/naloxone (Suboxone), the recommended first-line therapy. 16.9% were dispensed slow-release oral morphine (Kadian), while a small percent were prescribed the injectable buprenorphine-extended release (Sublocade) intended for moderate to severe opioid-use disorder management. Note that some clients might be dispensed more than one type of OAT in a given month.



Percentage of FNHA Clients Dispensed OAT for the First Time through the FNHA Health Benefits Plan by



Of all 2,557 FNHA clients dispensed OAT in May 2022, 2.5% were dispensed OAT through the FNHA health benefits plan for the first time.

Naloxone Distribution

Naloxone is an opioid antagonist that is used in an emergency response situation to temporarily reverse the effects of life-threatening opioid overdose. It is available in injectable or nasal spray form and often is bundled with other supplies (such as gloves or a breathing mask) in a carrying case or kit. The nasal spray is provided by the FNHA through two routes: by way of community pharmacies to First Nations individuals and through bulk supply to communities and Indigenous service organizations:

- Through FNHA's bulk ordering program, 1060 nasal naloxone kits were distributed to First Nations and community organizations in May 2022, and another 193 doses of nasal naloxone were distributed to First Nations clients through community pharmacies. FNHA Nasal Naloxone fact sheet
- Additionally, 720 injectable naloxone kits were ordered by 163 First Nations sites or Friendship Centres in May 2022. Injectable naloxone is available for free in the province to anyone at risk of an overdose or likely to witness one. For information on how to access and use an injectable naloxone kit, see Toward the Heart



Harm Reduction on FNHA.ca

For information about substance use, to get informed, and to support others, visit <u>Harm Reduction on</u> FNHA.ca, which includes:

- <u>Get Help</u>: Virtual Substance Use and Psychiatry Services; harm reduction services, including OPS/Raven's Eye Sites, naloxone (nasal naloxone and FNHA community bulk purchase), workshops including Not Just Naloxone, Decolonizing Substance Use, and Tackling Stigma, land-based healing programs, OAT, and drug testing
- <u>Get Informed</u>: personal stories about overdose and harm reduction; FNHA harm reduction campaign; learning resources; news; FNHA's Framework for Action; FNHA toxic drug annual data releases; and <u>Indigenous treatment centres</u>
- <u>Support Others</u>: Indigenous harm reduction; <u>Take-Home Naloxone</u> for the FNHA nasal naloxone programs; FNHA Indigenous Wellness Program; and learning resources for helping people who use substances

Latest News

- New Project Recognizes Grief Experienced by People Who Have Lost a Loved One to Toxic Drugs.
 October 3, 2022.
- On International Overdose Awareness Day 2022, We Honour Loved Ones Lost to the Toxic Drug Crisis and Intensify Our Actions. September 1, 2022.
- Reframing Indigenous Harm Reduction Ideologies to Heal Indigenous Spirit. August 31, 2022.
- Possession of Small Amounts of Drugs to Be Decriminalized but Not Legalized in BC. June 6, 2022.



Appendix: Data Sources and Definitions

BC Coroners Drug Toxicity Data

As defined by the BC Coroners Service (BCCS), "illicit drug overdoses include those involving street drugs (controlled and illegal: heroin, cocaine, MDMA, methamphetamine etc.), medications that were not prescribed to the deceased, combinations of the above with prescribed medications and those overdoses where the origin of the drug is not known. Both open and closed cases are included." (BCCS, 2018).

BCCS operates in a live database and includes both open and closed cases. Thus, data are subject to change as investigations are completed and data is refreshed. Small changes in numbers of deaths are expected with every refresh.

First Nations—specific information is identified via linkage to the FNCF, a cohort of all individuals registered with Indigenous Services Canada (ISC) as of 2018 and living in BC, as well as their eligible descendants. Only persons identified as status First Nations are captured via linkage. First Nations people without status, Métis and Inuit persons are not captured in the above data.

BC Emergency Health Services (BCEHS) Paramedic-Attended Drug Poisonings

Identification of drug-poisoning records is based on paramedic impression codes as well as 911 dispatch codes or where naloxone was administered by a paramedic. Alcohol and prescription drug related overdoses are excluded.

The majority of drug poisoning events identified by BCEHS data are nonfatal; however, it is possible that some deaths are also captured (BCCDC, 2021). Paramedic-attended toxic drug events include all events where 911 was called and BCEHS paramedics responded. Drug poisonings reversed in community where paramedics were not called are not captured.

Linkages to the FNCF requires a PHN. When a PHN is unavailable, the FNHA is not able to identify whether the record was of a First Nations persons or not. In 2021, approximately 25 per cent of events did not have a PHN; in 2020, approximately 24 per cent of events did not have a PHN; and in 2019, approximately 18 per cent of events did not have PHN and were thus not linkable to the FNCF. Consequently, paramedic-attended drug poisonings are likely underestimated for First Nations people. Additionally, there is likely a greater underestimation for 2020 and 2021 compared to previous years due to higher numbers of events in which a PHN was not available in these years. BCEHS is able to recover some of the missing PHNs; however, this process takes time. The Ministry of Health is able to run an additional algorithm to recover PHNs for some of the records. This absence of data is expected to decline with time.

First Nations data includes only persons with status and their descendants. First Nations persons without status are not included.

FNHA Health Benefits OAT Data

OAT data comes from line-level claims data for pharmacy dispensations through the First Nations Health Benefits program. There are three sources of this data: the federal Non-Insured Health Benefits (NIHB) program (up to September 15, 2019), BC PharmaCare Plan W (since September 2017), and Pacific Blue Cross Parallel Plan W (since September 2019). Since October 2017, 97.4% of FNHA clients have been enrolled in Plan W.

All measures in this report are broken down by provider region, except for unique prescriber counts which are broken down by assumed prescriber region.